

---

**INDICATIONS   CONTRA-INDICATIONS   DOSAGE   SIDE-EFFECTS   PREGNANCY**  
**OVERDOSE   IDENTIFICATION   PATIENT INFORMATION**

---

## TCP Antiseptic Liquid.

### SCHEDULING STATUS:

Unscheduled

### PROPRIETARY NAME

(and dosage form):

### TCP Antiseptic Liquid.

### COMPOSITION:

Each mL of TCP antiseptic contains, Chlorinated Phenols 6,0 mmg; Phenol 1,75 mg; Iodinated Phenols 0,95 mg; Sodium Salicylate 0,5 mg.

### PHARMACOLOGICAL CLASSIFICATION:

A 13.1 Antiseptics and Cleansing agents.

### PHARMACOLOGICAL ACTION:

TCP Antiseptic is a disinfectant effective against gram-positive and gram-negative bacteria.

### INDICATIONS:

TCP Antiseptic helps destroy bacteria and germs associated with throat and mouth infections. TCP Antiseptic by it's disinfectant action on bacteria and germs helps heal cuts, grazes, wounds, insect bites and burns on the skin.

TCP Antiseptic can be used as a personal hygiene, as a vaginal douche. TCP Antiseptic can be used as a general disinfectant wherever common germs are a danger to health around the home. TCP Antiseptic fights bad breath by acting on the odour-causing bacteria present in the mouth.

### CONTRA-INDICATIONS:

There are no known contraindications.

### DOSAGE AND DIRECTIONS FOR USE:

*Adults and Children*

#### MOUTH AND THROAT INFECTIONS

Gargle thoroughly with 1 part TCP Antiseptic to 4 or 5 parts water.

#### CUTS, GRAZES, WOUNDS, INSECT BITES AND BURNS.

First wash the affected area, then apply 1 part TCP Antiseptic to 2 or 3 parts water to thoroughly cleanse the area. If necessary fix lightly over the affected area a gauze damped with 1 part TCP Antiseptic to 2 or 3 parts water and hold in position with an adhesive plaster. Do not use heavy bandaging as wounds heal faster when air can reach them.

#### FEMININE HYGIENE

Douche with 4 tablespoons (75 mL) TCP Antiseptic to 500 mL warm water.

#### GENERAL HOUSEHOLD USE

TCP Antiseptic can be used undiluted on drains, sinks, dustbins and with the household laundry.

#### BAD BREATH

Rinse the mouth and gargle thoroughly with 1 part TCP Antiseptic to 4 or 5 parts water.

If infection persists consult your doctor immediately.

### SIDE EFFECTS AND SPECIAL PRECAUTIONS:

No side effects have been reported. TCP Antiseptic should not be drunk in large quantities as this may cause adverse reactions on the kidneys.

### KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

The treatment of overdosage is supportive and symptomatic.

### IDENTIFICATION:

TCP Antiseptic is a yellow clear liquid.

### PRESENTATION:

TCP Antiseptic is available in glass bottles of 200 mL, 100 mL and 50 mL individually packaged in carton boxes.

**STORAGE INSTRUCTIONS:**

Store below 30°C.

**Keep out of reach of children**

**REGISTRATION NUMBER:**

Applic. No. G1624 (Act 101/1965).

**APPLICANT:**

**Pfizer Laboratories (Pty) Limited**

102 Rivonia Road

Sandton, 2199

**DATE OF PUBLICATION:**

1-2-1982

---

[SAEPI HOME PAGE](#)

[TRADE NAME INDEX](#)

[GENERIC NAME INDEX](#)

[FEEDBACK](#)

---

Information presented by Malahyde Information Systems © Copyright 1996-2000